ISSN (print) 0093-4666 © 2014. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/127.1

Volume 127, pp. 1–4 *January–March 2014*

A new species of *Helminthosporium* from Jiangsu, China

Xiao-Jing Wang, Hai-Yan Wu, & Meng Zhang*

Henan Agricultural University Zhengzhou, Henan, China, 450002
* Correspondence to: *zm2006@126.com*

Abstract— *Helminthosporium nanjingense* sp. nov. is described from Jiangsu Province, China, on dead branches of an unidentified tree. The new fungus produces a conspicuous yellow-green pigment on PDA media. Its microscopic features and ITS sequence are provided.

Key words — systematics, saprobes, hyphomycetes

Introduction

During the recent survey of *Helminthosporium* species in China, the authors discovered eleven new species and seven new records (Zhang & Zhang 2009, Zhang et al. 2003, 2004, 2007, 2010). In this paper we propose a new species of *Helminthosporiu*m from Jiangsu Province, China.

Materials & methods

Unidentified dead branches with *Helminthosporium* species were carefully inspected under a Nikon SMZ1500 stereomicroscope. Tissues mounted in a drop of lactophenol were examined microscopically. Conidia and conidiophores were photographed using a Nikon Eclipse 80i microscope with Nikon DS-U2 camera, and microscopic measurements were completed by using NIS-Elements Image Analysis Software.

DNA was obtained from mycelia scraped from single spore cultures of the strains on a PDA plate and extracted using the Cell Genomic DNA Isolation Kit (LifeFeng, Shanghai, China) following the manufacturer's instructions. A final volume of 25 μL [comprising 1 μL ITS1 forward sequencing primer (5μM) and 1 μL ITS4 reverse sequencing primer (5 μM), 9.5 μL ddH_2O, 12.5 μL 2× Taq PCR Master Mix (including 0.1 U/μL Taq DNA Polymerase, 0.4 mM each dNTPs, 2× Taq Buffer, red and blue indicator dye, PCR enhance reagent and proteide steady reagent), 1 μL fungal DNA] was amplified using a Eppendorf PCR thermal cycler device (Germany) as follows: 95°C initial denaturation (2 min), 33cycles of 95°C (30 s), 55°C (30 s), 72°C (2 min), and a final step of 72°C (7 min). The 5-μL PCR products were analyzed through electrophoresis on a 1.0% agarose gel with 1× Tris-acetate-EDTA (TAE electrophoresis buffer). The PCR products were sequenced by Shanghai Sangon Biological Engineering Technology & Service Co. Ltd.,

China (Xie et al. 2010; Schoch et al. 2012). The type specimen is deposited in HSAUP (Herbarium of Shandong Agricultural University: Plant Pathology), and the ex-type culture in HHAUF (Herbarium of Henan Agricultural University: Fungi).

Taxonomy

Helminthosporium nanjingense Meng Zhang, Xiao J. Wang & H.Y. Wu, **sp. nov.**

Fig. 1

MycoBank MB 804655

Differs from other similar species of *Helminthosporium* by its smooth-walled, longer, slightly thinner conidia and its longer conidiophores.

Type: China, Jiangsu Province, Nanjing, on dead branches of an unidentified tree, 8 Sep. 2002, coll. M. Zhang (**Holotype**, $HSAUP_{02}0198$; ex-type culture, $HHAUF_{02}0380$=$ZM_{02}0380$; GenBank KF192322).

Etymology: Latin, *nanjingense,* for the collection locality.

Colony on the natural substrate effuse, black, hairy. Mycelium mostly immersed in the substratum. Colony on PDA media effuse with yellow-green pigment. Conidiophores arising solitary or in fascicles from the cells of the mycelium, simple, straight or flexuous, thick-walled, sub-cylindrical, smooth, brown to dark brown, 250–470 µm long, 6.9–7.7 µm wide, with well-defined small pores at the apex and laterally beneath the upper 1–4 septa; 8.6–11 µm wide at the base cell which sometimes inflated up to 15.5 µm. Conidia subulate or nearly whip-like, straight or curved, thin-walled, smooth, pale brown, 6–17 distoseptate, 64.5–170.5 µm long, 7.3–10.3 µm wide in the widest part, narrowing towards the apex to 5.0–6.8 µm wide, with a blackish-brown scar at the base, 1.4–2.7 µm thick.

Comments: *Helminthosporium nanjingense* is most closely related to *H. conidiophorellum* Meng Zhang & T.Y. Zhang and *H. subhyalinum* Meng Zhang & T.Y. Zhang in conidial morphology (subulate or nearly whip-like shape). *Helminthosporium conidiophorellum* differs from *H. nanjingense* by its shorter and slightly wider conidia (100–147.5 × 9.5–11 µm) that are sometimes slightly verruculose and its shorter wider conidiophores (60–280 × 7.0–8.5 µm) (Zhang & Zhang 2009). *Helminthosporium subhyalinum* is differentiated by its nearly colorless, shorter, slightly wider conidia (72–125 × 9.5–11.5 µm) and shorter conidiophores (120–200 µm); the thick black conidial apex wall of *H. subhyalinum* is also a striking distinguishing characteristic (Zhang et al. 2007).

The amplified rDNA (ITS) region derived from the holotype was 454 bp (GenBank KF192322). The BLAST comparison with previously published *H. solani* and *H. velutinum* sequences shows only 89% nucleotide homologies for both. Reference sequences for comparative analysis were obtained from



Fig. 1 *Helminthosporium nanjingense* (holotype HSAUP $_{02}$01980): A–D. Conidia; E. Conidiophore apex; F. Conidiophores; G. Conidiophore base. Scale bars = 50 μm.

GenBank database by the BLAST search program (http://www.ncbi.nlm.nih.gov/BLAST/).

Acknowledgments

We are deeply grateful to Drs. R.F. Castañeda Ruiz (Instituto de Investigaciones Fundamentales en Agricultura Tropical "Alejandro de Humboldt," Cuba) and Prof. Y.L. Guo (Institute of Microbiology, Academia Sinica, China) for kindly serving as pre-submission reviewers and their useful suggestions. This project was supported by the National Natural Science Foundation of China (Project No. 30970016, 30870018 & 31171804) and Sponsored by Program for Science & Technology Innovation Talents in Universities of Henan Province (No. 13HASTIT007).

Literature cited

Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, et al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. Proceedings of the National Academy of Sciences 109(16): 6241–6246.

Xie L, Huang SL, Cen ZL, Lu WH, Qin BX, Tang CG, Hu CJ, Qin LP. 2010. First report of *Botryosphaeria dothidea* causing sweet osmanthus leaf dieback in China. Agricultural Sciences in China 9(6): 847–853. http://dx.doi.org/10.1016/S1671-2927(09)60163-2

Zhang M, Zhang TY. 2009. Taxonomic studies of *Helminthosporium* from China 4. Six new species and a key to *Helminthosporium* from China. Mycotaxon 109: 399–413. http://dx.doi.org/10.5248/109.399

Zhang M, Zhang TY, Wu WP. 2003. Taxonomic studies of *Helminthosporium* from China I. Some new records. Mycosystema 22(Supplement): 77–79.

Zhang M, Zhang TY, Wu WP. 2004. Taxonomic studies of *Helminthosporium* from China II. Two new species in Sichuan Province. Mycosystema 23: 179–182.

Zhang M, Zhang TY, Wu WP. 2007. Taxonomic studies of *Helminthosporium* from China III. Three new species in Guangdong Province. Mycotaxon 99: 137–142.

Zhang M, Wu HY, Wang ZY. 2010. Taxonomic studies of *Helminthosporium* from China 5. Two new species from Hunan and Sichuan Province. Mycotaxon 113: 95–99. http://dx.doi.org/10.5248/113.95